

Study on two coupled relativistic superfluids with spontaneous symmetry breaking

Sun Zhang

Department of Physics, Nanjing University, Nanjing 210093, China

Abstract

In this Letter, we have studied two coupled relativistic superfluids with spontaneous U(1) symmetry breaking, using Poisson bracket technique. After constructing the commutators between thermo-quantities and field quantities, the equations of motion are obtained. These equations describe the system in the frame of two-constituent superfluid theory and provide a clear picture relating the symmetry breaking and the physical quantities explicitly. The interference effect is discussed and the Josephson-type equation is also given. Furthermore, the dissipation effect and entropy production mechanism are presented and the dissipative coefficients are given explicitly in the first-order theory framework.

PACS: 47.75.+f; 47.37.+q; 67.40.Bz

In 1941, Landau proposed his famous two-constituent theory on superfluid ^4He [1,2]. The central idea of the theory is that at temperature below the critical point, helium behaves as if it were a mixture of two different liquids. One of these is a superfluid, and moves with zero viscosity along a solid surface, while the other is a normal viscous liquid. No momentum is transferred from one to the other. The two-constituent superfluid theory was studied and generalized by Khalatnikov [3] brilliantly in the non-relativistic frame.

The first approach to the relativistic superfluid theory was provided by Israel [4] and Dixon [5,6] concerning the perfect fluids based on the idea of two-constituents originated by Landau. The medium is usually called relativistic in two senses: if it has a relativistic equation of state or when it flows at a relativistic velocity. While these conditions can be both satisfied in the neutron stars whose interior construction is composed of a superfluid nuclear matter [7]. The method proposed by Israel and Dixon is useful for some calculations, but arising a problem.

Generally speaking, the coupled two-constituents are not perfect fluids strictly and the superfluid component cannot be regarded as a completely independent fluid [8]. That is, the coupling effects lead the deviation from ideality [9]. It is meaningful to take into account the deviation from the perfect fluids. Khalatnikov and Lebedev [8] and Carter and Khalatnikov [10] suggested two equivalent approaches to include the interaction between the superfluid and normal fluid.

Furthermore, Son [11] presented another approach to the relativistic superfluid with $U(1)$ symmetry breaking based on the Poisson bracket technique [12]. Such a description is also arisen from the two-constituent theory, but it is more clearer to show the relation between superfluid and the symmetry breaking. After constructing some basic relations of the fundamental quantities by Poisson brackets, the dynamic equations are obtained directly, which provide the explicit meaning about the symmetry breaking.

It is possible to generalize the discussion presented in Ref. [11] in several directions. The most important one is that we should construct a clear frame to discuss the effects between two coupled relativistic superfluids including the coherence effect and the dissipation effect.

For the symmetry spontaneous breaking and the phase coherence are assumed from the very beginning, such an approach could be looked as a natural framework for investigating the inter-relationship of two coupled relativistic superfluids in the hydrodynamic limit.

To study the two coupled relativistic superfluids 1 and 2 with U(1) symmetry breaking, the order parameter is of the form $\langle\psi_i\rangle = |\langle\psi_i\rangle| e^{i\phi_i}$ ($i = 1, 2$), the density of the U(1) charge $\rho_i = -i(\psi_i^* \partial_0 \psi_i - \partial_0 \psi_i^* \psi_i)$ and the phase ϕ_i are introduced [11].

In order to establish the dynamic equations, the Poisson brackets should be written down to give the commutative relations between dynamic variables. There are two kinds of Poisson brackets: the fluid (thermo-) ones and the field ones.

The fluid ones are [11,12]

$$[T^{0i}(\mathbf{x}), A(\mathbf{y})] = A(\mathbf{x}) \partial_i \delta(\mathbf{x} - \mathbf{y}), \quad A = \rho_1, \rho_2, s \quad (1)$$

$$[T^{0i}(\mathbf{x}), B(\mathbf{y})] = -\partial_i B(\mathbf{x}) \delta(\mathbf{x} - \mathbf{y}), \quad B = \phi_1, \phi_2 \quad (2)$$

$$[T^{0i}(\mathbf{x}), T^{0j}(\mathbf{y})] = \left(T^{0j}(\mathbf{x}) \frac{\partial}{\partial x^i} - T^{0i}(\mathbf{y}) \frac{\partial}{\partial y^j} \right) \delta(\mathbf{x} - \mathbf{y}), \quad (3)$$

where T^{0i} is the momentum density in the 4-dimensional energy-momentum tensor, s the entropy density. $\rho_{1,2}$ and $\varphi_{1,2}$ are just the density of charge and the phase of the order parameter respectively.

The Poisson bracket for the field part includes the commutator of ρ_i with ϕ_j (up to an adjustable factor)

$$[\rho_i(\mathbf{x}), \phi_j(\mathbf{y})] = -\delta(\mathbf{x} - \mathbf{y}) \delta_{ij}. \quad i, j = 1, 2 \quad (4)$$

To get the equation of motion, we should construct the most general Hamiltonian dependent on the fluid quantities and the field quantities

$$H = \int d\mathbf{r} T^{00}(s, \rho_1, \rho_2, T^{0i}, \partial_i \phi_1, \partial_i \phi_2), \quad (5)$$

where T^{00} is a functional of $\partial_i \phi_{1,2}$ but not $\phi_{1,2}$ itself due to the U(1) invariance [13,14]. The form of the T^{00} cannot be identified at this moment, it could be calculated from the microscopic theory.

Then the equation of state, describing the relation of the pressure p , the temperature T , the chemical potentials $\mu_{1,2}$, and the phases $\phi_{1,2}$, could be written in the differential form as

$$dp = sdT + \rho_1 d\mu_1 + \rho_2 d\mu_2 + V_{11}^2 d\left(\frac{1}{2}(\partial_\mu \phi_1)^2\right) + V_{22}^2 d\left(\frac{1}{2}(\partial_\mu \phi_2)^2\right) - (V_{12}^2 + V_{21}^2) d\left(\frac{1}{2}\partial_\mu \phi_1 \partial^\mu \phi_2\right), \quad (6)$$

where, V_{11}^2 and V_{22}^2 are the factors characterizing the strength of the internal energy for superfluids 1 and 2 respectively, while V_{12}^2 and V_{21}^2 are the coupling constants between two superfluids due to interference and $V_{12}^2 = V_{21}^2$. In Eq. (6), the first three terms are fluid terms, they describe the normal fluids part, while the gradients of the phases in the left terms are specific to the superfluids and have the significance of the superfluid velocities.

Based on the Poisson brackets (1)–(4) and the thermodynamic relations, we obtain the dynamic equations

$$\partial_\mu \left(u^\mu \rho_1 - V_{11}^2 \partial^\mu \phi_1 + \frac{1}{2}(V_{12}^2 + V_{21}^2) \partial^\mu \phi_2 \right) = 0, \quad (7)$$

$$\partial_\mu \left(u^\mu \rho_2 - V_{22}^2 \partial^\mu \phi_2 + \frac{1}{2}(V_{12}^2 + V_{21}^2) \partial^\mu \phi_1 \right) = 0, \quad (8)$$

$$u^\mu \partial_\mu \phi_1 + \mu_1 = 0, \quad (9)$$

$$u^\mu \partial_\mu \phi_2 + \mu_2 = 0. \quad (10)$$

It is clear that Eqs. (7) and (8) are the conservation laws of the U(1) charge in the Lorentz covariance form. Within the bracket of Eq. (7) [or Eq. (8)], the first term presents the normal current and the 4-dimensional velocity u^μ is the velocity of the normal component; while the second term gives the superfluid current and the gradient of the phase is clearly the velocity of the superfluid component, and the two-constituent description is constructed explicitly. The third term is the interference current, which illuminates the phase coherent influence of the superfluids from each other. Then the U(1) charge is a sum of the normal

current, the superfluid current, and the interference current due to the coupling to the other superfluid, and Eqs. (7) and (8) give the conservation relations of the U(1) charge.

On the other hand, Eq. (9) [or Eq. (10)] gives the time dependence of the condensate phase on the chemical potential. The chemical potential in these equations is just the change in energy density per change in the charge density, and also the minimum energy required to add a particle to the system. The chemical potential also has the meaning of the background field due to the U(1) gauge transformation, and coincides the scalar potential V in the superconductivity. If we suppose that the two superfluids are kept at uniform chemical potentials, it is direct to obtain the time dependent relation of the difference of the condensate phases according to Eqs. (9) and (10)

$$u^\mu \partial_\mu (\phi_1 - \phi_2) + (\mu_1 - \mu_2) = 0. \quad (11)$$

It is just the relativistic version of the Josephson equation, in BEC coherence effect [15] and superconductivity [16].

We can also construct the energy-momentum tensor $T^{\mu\nu}$ as a sum of the thermo (fluid) part and field (superfluid) part as follows

$$T^{\mu\nu} = (\epsilon + p)u^\mu u^\nu - pg^{\mu\nu} + V_{11}^2 \partial^\mu \phi_1 \partial^\nu \phi_1 + V_{22}^2 \partial^\mu \phi_2 \partial^\nu \phi_2 - V_{12}^2 \partial^\mu \phi_1 \partial^\nu \phi_2 - V_{21}^2 \partial^\mu \phi_2 \partial^\nu \phi_1, \quad (12)$$

which satisfies the conservation law

$$\partial_\mu T^{\mu\nu} = 0, \quad (13)$$

where the fluid part [2]

$$T_{\text{fluid}}^{\mu\nu} = (\epsilon + p)u^\mu u^\nu - pg^{\mu\nu}, \quad (14)$$

describes the normal relativistic fluid outside the superfluid condensation phase and the metric tensor is

$$g^{\mu\nu} = \text{diag}(1, -1, -1, -1), \quad (15)$$

in accordance with Landau and Lifshitz, and $u^\mu u_\mu = 1$. While the field part describes the coherent motion and the interference effect of two superfluids.

Next, we would like to explore the dissipation effect and the entropy production mechanism of two coupled relativistic superfluids system to provide a full understanding of the behaviors described here, following Landau and Lifshitz [2] and Weinberg [17].

Generally speaking, the dissipative processes are important, when space-time gradients of hydrodynamic quantities in the system become large relative to its relaxation scales. We would analyze the dynamics of small departures of the system from equilibrium states and manage to give the dissipative equations as linear responses to the deviations from equilibrium states and two famous dissipative theories by Eckart [18] and by Landau and Lifshitz [2] would be the special cases in the theory presented here.

When we study the imperfect system of two coupled superfluids, the dissipative effects should be taken into account by modifying the energy-momentum tensor $T^{\mu\nu}$ by a term $\Delta T^{\mu\nu}$, and could be written in matrix form as

$$T^{\mu\nu} + \Delta T^{\mu\nu} = (\epsilon + p)u^\mu u^\nu - pg^{\mu\nu} + \begin{pmatrix} \partial^\mu \phi_1 & \partial^\mu \phi_2 \end{pmatrix} \begin{pmatrix} V_{11}^2 & -V_{12}^2 \\ -V_{21}^2 & V_{22}^2 \end{pmatrix} \begin{pmatrix} \partial^\nu \phi_1 \\ \partial^\nu \phi_2 \end{pmatrix} + \Delta T^{\mu\nu}, \quad (16)$$

satisfying the conservation equation as

$$\partial_\mu (T^{\mu\nu} + \Delta T^{\mu\nu}) = 0. \quad (17)$$

The other dissipative effects for the charge and the phase could be considered as

$$\partial_\mu \begin{pmatrix} u^\mu \rho_1 + \Delta \rho_1^\mu \\ u^\mu \rho_2 + \Delta \rho_2^\mu \end{pmatrix} = \partial_\mu \left[\begin{pmatrix} V_{11}^2 & -V_{12}^2 \\ -V_{21}^2 & V_{22}^2 \end{pmatrix} \begin{pmatrix} \partial^\mu \phi_1 \\ \partial^\mu \phi_2 \end{pmatrix} \right], \quad (18)$$

and

$$u^\mu \partial_\mu \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} = - \begin{pmatrix} \mu_1 + \Delta \phi_1 \\ \mu_2 + \Delta \phi_2 \end{pmatrix}. \quad (19)$$

The dissipative terms should satisfy the following constraints

$$u_\mu u_\nu \Delta T^{\mu\nu} = 0, \quad (20)$$

$$u_\mu \Delta \rho_{1,2}^\mu = 0. \quad (21)$$

For comparison, the constraint used by Landau and Lifshitz is $u_\nu \Delta T^{\mu\nu} = 0$ and that of Eckart is $\Delta \rho^\mu = 0$.

Then using the equations of motion and the thermodynamic relations, we can get the entropy production equation arisen by the dissipative terms,

$$\begin{aligned} \partial_\mu S^\mu = & -\partial_\mu \left(\frac{\mu_1}{T}, \frac{\mu_2}{T} \right) \begin{pmatrix} \Delta \rho_1^\mu \\ \Delta \rho_2^\mu \end{pmatrix} \\ & + \left(\frac{\Delta \phi_1}{T}, \frac{\Delta \phi_2}{T} \right) \partial_\mu \left[\begin{pmatrix} V_{11}^2 & -V_{12}^2 \\ -V_{21}^2 & V_{22}^2 \end{pmatrix} \begin{pmatrix} \partial^\mu \phi_1 \\ \partial^\mu \phi_2 \end{pmatrix} \right] \\ & + \frac{1}{T} \Delta T^{\mu\nu} \partial_\mu u_\nu - \frac{1}{T^2} u_\mu \Delta T^{\mu\nu} \partial_\nu T. \end{aligned} \quad (22)$$

Where the entropy S^μ density is defined as

$$S^\mu = u^\mu s - \left(\frac{\mu_1}{T}, \frac{\mu_2}{T} \right) \begin{pmatrix} \Delta \rho_1^\mu \\ \Delta \rho_2^\mu \end{pmatrix} + \frac{1}{T} u_\nu \Delta T^{\mu\nu}. \quad (23)$$

By means of the projection operator $(g^{\mu\nu} - u^\mu u^\nu)$, $\Delta T^{\mu\nu}$ can be expressed as

$$\begin{aligned} \Delta T^{\mu\nu} = & ((g^{\mu\alpha} - u^\mu u^\alpha)u^\nu + (g^{\nu\alpha} - u^\nu u^\alpha)u^\mu) q_\alpha \\ & + (g^{\mu\alpha} - u^\mu u^\alpha)(g^{\nu\beta} - u^\nu u^\beta) \tau_{\alpha\beta} + (g^{\mu\nu} - u^\mu u^\nu) \tau. \end{aligned} \quad (24)$$

Defining the quantities $\Delta N_{1,2}$ as

$$\begin{pmatrix} \Delta N_1 \\ \Delta N_2 \end{pmatrix} = \begin{pmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{pmatrix} \begin{pmatrix} \frac{\mu_1}{T} \\ \frac{\mu_2}{T} \end{pmatrix}, \quad (25)$$

where σ_{11} and σ_{22} are the charge diffusion coefficients for superfluids 1 and 2 respectively, while $\sigma_{12} = \sigma_{21}$ are the mutual diffusion coefficients between two superfluids and are supposed to have only nonnegative eigenvalues. Then the charge dissipative equations are

$$\begin{aligned}
\begin{pmatrix} \Delta\rho_1^\mu \\ \Delta\rho_2^\mu \end{pmatrix} &= (g^{\mu\nu} - u^\mu u^\nu) \partial_\nu \left[\begin{pmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{pmatrix} \begin{pmatrix} \frac{\mu_1}{T} \\ \frac{\mu_2}{T} \end{pmatrix} \right] \\
&= (g^{\mu\nu} - u^\mu u^\nu) \partial_\nu \begin{pmatrix} \Delta N_1 \\ \Delta N_2 \end{pmatrix}.
\end{aligned} \tag{26}$$

Then the first term in Eq. (22) could be written as

$$\begin{aligned}
-\partial_\mu \left(\frac{\mu_1}{T}, \frac{\mu_2}{T} \right) \begin{pmatrix} \Delta\rho_1^\mu \\ \Delta\rho_2^\mu \end{pmatrix} &= -(g^{\mu\nu} - u^\mu u^\nu) \partial_\mu \left(\frac{\mu_1}{T}, \frac{\mu_2}{T} \right) \partial_\nu \begin{pmatrix} \Delta N_1 \\ \Delta N_2 \end{pmatrix} \\
&= -(g^{\mu\nu} - u^\mu u^\nu) \partial_\mu (\Delta N_1, \Delta N_2) \begin{pmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{pmatrix}^{-1} \partial_\nu \begin{pmatrix} \Delta N_1 \\ \Delta N_2 \end{pmatrix},
\end{aligned} \tag{27}$$

where the positivity of the inverse matrix of σ is assumed. Defining the quantities $A_{1,2}^\mu$ as

$$\begin{pmatrix} A_1^\mu \\ A_2^\mu \end{pmatrix} = \begin{pmatrix} V_{11}^2 & -V_{12}^2 \\ -V_{21}^2 & V_{22}^2 \end{pmatrix} \begin{pmatrix} \partial^\mu \phi_1 \\ \partial^\mu \phi_2 \end{pmatrix}, \tag{28}$$

and $\Delta M_{1,2}^\mu$ as

$$\begin{pmatrix} \Delta M_1^\mu \\ \Delta M_2^\mu \end{pmatrix} = \begin{pmatrix} \zeta_{11} & \zeta_{12} \\ \zeta_{21} & \zeta_{22} \end{pmatrix} \begin{pmatrix} A_1^\mu \\ A_2^\mu \end{pmatrix}, \tag{29}$$

where ζ_{11} and ζ_{22} are the bulk viscosity coefficients for superfluids 1 and 2 respectively, while $\zeta_{12} = \zeta_{21}$ are the mutual viscosity coefficients between two superfluids and have only nonnegative eigenvalues supposedly.

Then the second term in Eq. (22) could be written as

$$\begin{aligned}
&\left(\frac{\Delta\phi_1}{T}, \frac{\Delta\phi_2}{T} \right) \partial_\mu \left[\begin{pmatrix} V_{11}^2 & -V_{12}^2 \\ -V_{21}^2 & V_{22}^2 \end{pmatrix} \begin{pmatrix} \partial^\mu \phi_1 \\ \partial^\mu \phi_2 \end{pmatrix} \right] \\
&= \left(\frac{\Delta\phi_1}{T}, \frac{\Delta\phi_2}{T} \right) \partial_\mu \begin{pmatrix} A_1^\mu \\ A_2^\mu \end{pmatrix} = \frac{1}{T} \partial_\mu (\Delta M_1^\mu, \Delta M_2^\mu) \begin{pmatrix} \zeta_{11} & \zeta_{12} \\ \zeta_{21} & \zeta_{22} \end{pmatrix}^{-1} \partial_\nu \begin{pmatrix} \Delta M_1^\nu \\ \Delta M_2^\nu \end{pmatrix} \\
&\quad + \frac{1}{T} \partial_\mu u^\mu (\zeta_1, \zeta_2) \partial_\nu \begin{pmatrix} A_1^\nu \\ A_2^\nu \end{pmatrix},
\end{aligned} \tag{30}$$

if we present the dissipative equation $\Delta\phi_{1,2}$ as

$$\begin{pmatrix} \Delta\phi_1 \\ \Delta\phi_2 \end{pmatrix} = \begin{pmatrix} \zeta_{11} & \zeta_{12} \\ \zeta_{21} & \zeta_{22} \end{pmatrix} \partial_\mu \begin{pmatrix} A_1^\mu \\ A_2^\mu \end{pmatrix} + \partial_\mu u^\mu \begin{pmatrix} \zeta_1 \\ \zeta_2 \end{pmatrix}, \quad (31)$$

where the viscosity coefficients ζ_1 and ζ_2 are those related to the normal velocity of the systems respectively. The dissipative components in $\Delta T^{\mu\nu}$ of Eq. (24) could be shown as

$$q_\alpha = \kappa T \left(\frac{1}{T} \partial_\alpha T - u^\mu \partial_\mu u_\alpha \right), \quad (32)$$

$$\tau_{\alpha\beta} = \eta (\partial_\alpha u_\beta + \partial_\beta u_\alpha - \frac{2}{3} g_{\alpha\beta} \partial_\mu u^\mu), \quad (33)$$

$$\tau = \zeta \partial_\mu u^\mu + (\zeta'_1, \zeta'_2) \partial_\mu \begin{pmatrix} A_1^\mu \\ A_2^\mu \end{pmatrix}, \quad (34)$$

where σ is the charge-diffusion coefficient, ζ is the total bulk viscosity coefficient, while κ and η are the thermal conductivity and the shear viscosity respectively. According to the Onsager's principle,

$$\zeta_i = \zeta'_i, \quad i = 1, 2 \quad (35)$$

and the positivity should be ensured by the following inequality

$$\zeta(\zeta_{11}\zeta_{22} - \zeta_{12}\zeta_{21}) \geq \zeta_2^2\zeta_{11} - 2\zeta_1\zeta_2\zeta_{12} + \zeta_1^2\zeta_{22}, \quad (36)$$

such an inequality is a generalization of that by Landau and Lifshitz [2].

Then the full set of dissipative equations are established here, and all transport coefficients are defined. These coefficients could be calculated from the microscopic theory and Kubo's linear response theory could be used here to build up the connections between the microscopic fluctuation and the macroscopic dissipation.

In summary, the fundamental equations of motion have been constructed which could be used to study the effects between two coupled relativistic superfluids. The theoretical

frame presented here are essentially based on the idea of two-constituent model proposed by Landau. Furthermore, in such a theoretical description, there arises more clearer physical significance for the equations of motion, which relate the symmetry breaking and the physical quantities naturally. Based on these, the interference effect and the Josephson-type equation are discussed; the dissipation effect and entropy production mechanism are presented, and the dissipative coefficients are given explicitly in the first-order theory framework.

Besides the interference effect between two coupled relativistic superfluids with spontaneous $U(1)$ symmetry breaking, another important effect is the discontinuity, including the shock wave propagation in the two-constituent relativistic superfluids. Although the shock wave was studied by many authors in the relativistic systems [19], a new theoretical frame is still necessary to provide some new features qualitatively. Instead of working in the frame of perfect fluid, we would rather select the theoretical picture presented here to give a new description on the shock wave in relativistic superfluids [20].

REFERENCES

- [1] L.D. Landau, J. Phys. USSR 5 (1941) 71.
- [2] L.D. Landau, E.M. Lifshitz, Fluid Mechanics, 2nd ed., Pergamon Press, Oxford, 1987.
- [3] I.M. Khalatnikov, An Introduction to the Theory of Superfluidity, Addison-Wesley, Redwood City, 1989.
- [4] W. Israel, Ann. Phys. (New York) 100 (1976) 310; Phys. Lett. A 86 (1981) 79.
- [5] W.G. Dixon, Thermodynamics of Superfluids via Relativity, Churchill College, Cambridge, 1981.
- [6] W.G. Dixon, Arch. Rat. Mech. Anal. 80 (1982) 159.
- [7] S. Shapiro, S. Teukolsky, Black Hole, White Dwarfs, and Neutron Stars, John Wiley and Sons, New York, 1983.
- [8] I.M. Khalatnikov, V.V. Lebedev, Phys. Lett. A 91 (1982) 70.
- [9] G.V. Vlasov, Sov. JETP 84 (1997) 729; hep-ph/9808250; hep-ph/0008219.
- [10] B. Carter, I.M. Khalatnikov, Ann. Phys. (New York) 219 (1992) 243; Phys. Rev. D 45 (1992) 4536.
- [11] D.T. Son, Int. J. Mod. Phys. A 16S1C (2001) 1284; hep-ph/0011246.
- [12] D.T. Son, Phys. Rev. Lett. 84 (2000) 3771; T.E. Dzyaloshinskii, G.E. Volovick, Ann. Phys. (New York) 125 (1980) 67.
- [13] D.T. Son, hep-ph/0204199.
- [14] S. Weinberg, The Quantum Theory of Fields, Vol. II, Cambridge University Press, Cambridge, England, 1996, Chapter 21.
- [15] F. Dalfovo, L. Pitaevskii, S. Stringari, Phys. Rev. A 54 (1996) 4213.

- [16] R.P. Feynman, Statistical Mechanics, Addison-Wesley, Redwood City, 1972.
- [17] S. Weinberg, Gravitation and Cosmology, Wiley, New York, 1972, Chapter 2.
- [18] C. Eckart, Phys. Rev. 58 (1940) 919.
- [19] A.H. Taub, Phys. Rev. 74 (1948) 328; M. Cissoko, Phys. Rev. D 55 (1997) 4555; K.A. Bugaev, M.I. Gorenstein, W. Greiner, Heavy Ion Phys. 10 (1999) 333.
- [20] S. Zhang, to be submitted.